

A note on magnetic-field induced level-density condensation in a two-dimensional electron gas with point scatterers

N. Kumar and Nivedita Deo

Raman Research Institute, C V Raman Avenue, Bangalore 560080, India

Abstract

The density-of-states (DOS) for a magnetized (B) two-dimensional electron gas (2DEG) containing point scatterers of arbitrary strengths, concentration (n_s) and distribution is analyzed. It is shown from the first principles that for $n_s \leq B/\Phi_o \equiv n_B$, the areal density of flux quanta $\Phi_o \equiv hc/e$, the DOS retains the extensive degeneracy characteristic of the Landau levels, but reduced by a factor $(1 - n_s/n_B)$. This elementary but exact result gives a level condensation for magnetic field $B > n_s\Phi_o$, as first noted by Brézin *et al.*. Its implications for the Integral Quantum Hall Effect and for Random Matrix Theory are pointed out.

PACS: 72.15 Gd, 71.25 Pi, 71.50 + t

Keywords: Integral Quantum Hall Effect, Disorder

Corresponding author: Prof. N.Kumar
Raman Research Institute
Bangalore 560080, India
Telephone: (+91 80) 331 1012
Fax: (+91 80) 334 0492
e-mail: nkumar@rri.ernet.in

A two-dimensional electron system in an external magnetic field (B) applied perpendicular to the plane, and with randomly distributed and strengthened impurities (disorder) is one of the most extensively studied low-dimensional systems, known best for the robust phenomenon of Integral Quantum Hall Effect (IQHE) [1]. The simplest model here consists of a two-dimensional electron gas (2DEG), i.e., without electron-electron interaction, spin-polarized in a strong uniform perpendicular magnetic field with uncorrelated point-like scatterers having a Poissonian distribution in the plane. It is described by the one-electron Hamiltonian [2]

$$H = \frac{1}{2m} \left(-i\hbar\nabla - \frac{e}{c}A \right)^2 + \sum_{i=1}^{N_s} V_i \delta(r - R_i) \equiv H_o + V \quad (1)$$

with the vector potential $\mathbf{A} = B(0, -x)$ in the Landau gauge. In the following we consider the general case of arbitrary spatial positions ($\{R_i\}$) and strengths ($\{V_i\}$) of the point scatterers. The mean concentration of scatterers $n_s = N_s/A$, with the area $A = L_x L_y$. Thermodynamic limit $L_x, L_y \rightarrow \infty$ will be assumed throughout.

The Hamiltonian H_o describes the well known Landau problem having eigenfunctions $\Phi_{nk}(\mathbf{r})$ and the corresponding eigenvalues $E_n = (n + \frac{1}{2})\hbar w_c$, labelled by the Landau-level index $n = 0, 1, 2, \dots$, and the quasi-continuous momentum label k taking on $N_B (= eB/hc)A$ values. Here $w_c = eB/mc$, the cyclotron frequency. Each Landau level is extensively degenerate with an areal density $n_B = B/\Phi_o$, where $\Phi_o = hc/e$ is the magnetic flux quantum.

The unperturbed DOS is thus given by

$$N_o(E; B) = (B/\Phi_o) \sum_{n=0}^{\infty} \delta(E - \hbar w_c(n + \frac{1}{2})) \quad (2)$$

We will now show that the exact DOS $N(E; B)$ for the disordered system continues to have an extensive degeneracy for $B > B_c = n_s \Phi_o$, characteristic of the Landau system, but reduced by a factor $(1 - n_s/n_B)$. This implies level condensation as the field exceeds the critical field B_c . The uncondensed fraction n_s/n_B forms a diffuse background to which the Wigner semi-circular law of Random Matrix Theory (RMT) [3] should apply.

With $N_B > N_s$, consider the n^{th} Landau level of H_o and construct the N_B degenerate linear combinations

$$\psi_n^m(\mathbf{r}) = \sum_{k=1}^{N_B} C_{nk}^m \phi_{nk}(\mathbf{r}), \quad m = 1, 2, \dots, N_B \quad (3)$$

as the new bases. Now, choose the coefficients $\{C_{nk}^m\}$ such that $\psi_n^m(\mathbf{r}) = 0$ for $r = R_i$, i.e., $\psi_n^m(\mathbf{r})$ has nodes at the sites of the point scatterers,

$$\psi_n^m(r = R_i) \equiv \sum_{k=1}^{N_B} C_{nk}^m \phi_{nk}(r = R_i) = 0 \quad (4)$$

for $m = 1, 2, \dots, N_s$.

Inasmuch as there are N_s conditions (equations) and N_B unknown coefficients, we are left with a sub-space of dimension $(N_B - N_s)$ spanned by the linear combinations ψ_n^m that vanish at $\{R_i\}$. Further, since these linear combinations vanish at the impurity sites $\{R_i\}$, they remain unaffected by the point scatterers [4]. Thus, there will be degenerate $(N_B - N_s)$ eigenstates (per Landau level) unaffected by the disorder. Accordingly, we have

the perturbed DOS

$$N(E; B) = \sum_n \left(1 - \frac{n_s}{n_B}\right) \delta(E - E_n) + N_{diffuse}(E) . \quad (5)$$

Thus, there is a condensation of level-density for a strong magnetic field. The condensation vanishes for $B \leq B_c = n_s \Phi_o$. The uncondensed fraction distributed as a band $N_{diffuse}(E)$ corresponds to the states affected by disorder. For a clean sample with $n_s \approx 10^{11} \text{ cm}^{-2}$, we have the critical field $B_c \approx 20\text{kG}$. The condensed eigenstates (extended) at the centre(s) of the Landau band(s) are believed to be important for IQHE [5].

It is now apt to compare the DOS in Eq.(5) to the one obtained perturbatively by Brézin et al. [2] (their Eq.50a) using the supersymmetric formalism. The condensate part matches exactly. However, it must be noted that their result was obtained specifically for the Poissonian distribution after averaging over the scattering impurity strengths, and for the single lowest Landau level (LLL). The present first-principle result for the condensation holds for the most general distribution of point scatters, and, in fact, without restriction to LLL. Physically, point scatterers would imply the range of scattering potential much smaller than the magnetic length ($= \sqrt{\hbar c/eB}$). For the diffuse part of DOS, however, we have no exact results here. But, the following argument should hold in the strong-field limit – if the inter-Landau level scattering is neglected as in ref. 3. (Though this may not be a very good approximation for the case of *abrupt*, short-ranged scatterers [6].) As the uncondensed fraction n_s/n_B of the total states in the LLL is subject to random scattering, one may expect the results of the

RMT to hold, and hence, in particular, we expect [3]

$$\frac{N_{diffuse}^{LLL}}{N_B}(E) = \frac{n_s}{2\pi\Delta n_B} \sqrt{4 - \left(\frac{E - E_o}{\Delta}\right)^2} \quad (6)$$

We would like to emphasize, however, that while the condensate part of the density of states Eq.(5) is exact for the point scatterer model, the diffuse part $N_{diffuse}$ is some non-universal background. Only for Gaussian RMT it assumes the semicircular form. We note here that a DOS with a delta-function and a semi-circular background has been obtained earlier [9] using RMT. In point of fact, in the absence of inter-Landau level scattering, all eigenstates can be readily shown to be extended [6,7] for the quantum-Hall-effect systems. And the extended states are, of course, a necessary condition for the RMT to hold. We can therefore treat Eqs.(5) & (6) as providing the input density of states for constructing the effective potential for the RMT unitary ensemble with the parameter (n_s/n_B) tunable by varying the magnetic field. We can then invoke the known expressions for the various correlation functions evaluated by Nagao *et al.* [10]. This then provides us with the possibility of studying these correlation functions as function of magnetic field for the quantum Hall regime. In particular, the coefficient ‘a’ (strength of the log potential in their Eqs. (2.2) & (2.3)) must vanish and stay zero for $B < B_c$. To our knowledge, this is the first time that a condensed matter system has been identified for the RMT-model potential giving the semi-circle plus delta function DOS. The non-condensed diffuse DOS obtained by Brézine *et al.* is, however, qualitatively different from the above semi-circular law.

We would like to conclude with a remark of direct relevance to the Hall conductance plateaus in the context of the above model (i.e., within the lowest Landau Level subspace). Normally, a Hall plateau represents the gap between the two mobility edges separating the adjacent Landau levels. As the disorder increases, the mobility edges are expected to move inward towards the respective Landau level centres, and hence the width of the plateau should increase, making the Hall steps steeper. In the present model, however, the opposite should happen. As all the states are extended, the plateaus can arise only due to the DOS gaps between the adjacent Landau levels [7]. Therefore, increasing the disorder strength would decrease the plateau-width as the DOS gap decreases because of the increasing width of the diffuse band ($N_{diffuse}(E)$) about the Landau levels. To the extent that the model of Brézinet *al.* is a good approximation in the high-field limit, the above opposite behaviour is inescapable.

Acknowledgements

N.D. would like to thank E. Brézin, R.N.Bhatt, F. David and S. Jain for discussions.

References

- [1] For a review, see R.E.Prange and S.M.Girvin, in The Quantum Hall Effect, 2nd Edn. (Springer, Berlin, 1990), and references cited therein.
- [2] E.Brézin, D.J.Gross and C.Itzykson, Nucl. Phys. B235 (1984) 24.
- [3] See M.L.Mehta, Random Matrices and the Statistical Theory of Energy Levels, 2nd Edn. (Academic, New York, 1991).
- [4] The point that the wavefunction and the eigenvalue remain unaffected by a point scatterer located at the node of the wavefunction has been noted by C.Itzykson in a preprint (1984). We are thankful to Prof. E. Brézin for bringing this to our attention.
- [5] See Y. Huo and R.N.Bhatt, Phys. Rev. Lett. 68 (1992) 1375 and references cited therein.
- [6] Robert Joynt and R.E.Prange, Phys. Rev. B 29 (1984) 3303.
- [7] N.Kumar, preprint (1996) cond-mat. # 9607064.
- [8] J.Bellissard, A. van Elst, and H.Schulz-Baldes, J. Math. Phys. 35 (1994) 5373.
- [9] C.-I.Tan, Mod. Phys. Lett. A 6 (1991) and Phys. Rev. D 45 (1992) 2862.
- [10] T. Nagao and K. Slevin, J. Math. Phys. 34 (1993) 2075.